Chapter 6: References

1. A.Kocisova, P.Novak, J. toporcak, M.Petrovsky,(2002), Development of Resistance in Field Housfly (*Musca domestica*): Comaparison of Effects of Classic Spray Regimes versus Integrated Control Methods., ACTA VET.BRNO, 71:401-405.

2. Ahmed Mohammed and Craig J. Coates,(2001), Towards germ line transformation of the potato tuber moth, Phthorimaea operculella (Zeller). Texas A&M University, Department of Entomology, Minnie Belle Heep Center, College Station, TX. Monday, December 10, - D0021.

3.Areekul, S., P. Sinchaisri and S. Tigvatananon, (1988). Effects of Thai plant extracts on the oriental fruit fly III.Attractancy Test. J. (Nat.Sci.) Kasetsart. 22(2): 160-164.

4. Arthur Woods,(1974), Pest Control: A survey, McGraw Hill Book Company.

5. Brader, L. (1979): Integrated pest control in developing world. *Ann. Rev. of Entomol*., 24, 225-254.

6. Brown,A.W.A.(1983): Insecticide resistance as a factor in the integrated control of Culicidae. In 'Integrated Mosquito Control Methodologies',Laird,M.,Miles,J.W.,Eds., Academic Press :New York, pp 161-235.

7. Carson R. (1962): Silent Spring, Honghton Mitfilin, Boston, Hamish Hamilton, London,1963.

8. Edwards C.A. (1973): Environmental pollution by pesticides, Plenum Press.

9.Ehrlich P.R. and Raven A.H. (1964): Evolution of butterflies and plants: A study in coevolution, *Evolution*,18, 586-608.

10. Environmental Protection Agency (EPA) (1993). Integrated Risk. Information System, d-Limonene. CASRN 5989-27-5.

11. Georghiou G.P. and Taylor C.E. (1986): Factors influencing the evolution of resistance. In 'National Research Council (U.S) Committee on Strategies for Management of Pesticide Resistant Pest Population. Ed. Pesticide Resistance: Strategies and tactics for management. National Academy press Washington D.C. pp 157-169.

12.J. Cameron[1], A.M. Gatland[2], G.P. Walker[1] and P.J. Wigley[2],(1997), Alternative Host Plants As Refugia For Diamondback Moth And Potato Tuber moth, [1]NZ Institute for Crop & Food Research Ltd, Private Bag 92169, Auckland [2]BioDiscovery NZ Ltd, 197 Atkinson Rd, Titirangi, Auckland.

13.Jaruwichitratana, S., P. Tanyasititsunthorn and P.Timpatanapong,(1988). Effect of citronella cream as a mosquito- repellent. Ramatibhadee; 11: 94-7.

14.Kanat; Amla,(2004); Insecticidal effect of essential oils from various plants against larvae of pine processionary moth,vol.60,no.2,pp.173-177(5).

15.Klocks J.A. (1987): Natural plant compounds, useful in insect control.*In allelochemicals: Role in Agriculture and Forestry*. G.R.Waller Ed.ACS Symposium ser.330, American Chemical Scoiety, Washington D.C. pp 396-415.

16.Law J.H.and Regnier F.E.(1971): Pheromones, *Ann. Rev. Biochem*., 40, 533.

17.M. HAFEZ, F. N. ZAKI, A. MOURSY, M. SABBOUR,(1994).Biological Effects of the Entomopathogenic Fungus, Beauveria Bassiana on the Potato Tuber Moth Phthorimaea Operculella (Seller) Journal of Islamic Academy of Sciences, Volume 7, No.4.Back to Journal of IAS, Volume 7, No 4.

18.Narongchai, P., S. Narongchai, S. Thampituk (2005). The First Fatal Case of Yam Bean and Rotenone Toxicity in Thailand. J Med Assoc Thai., 88(7): 984-986.

19.Norris D.M. and Kogan M. (1980): Biochemical and morphological basis of resistance. In 'Breeding plant resistant to insects'. Eds.E.G.Maxwell and P.R.Jenings. John Wiley and Sons, New York. 23-61.

20. Papachristos D. P. and D. C. Stamopoulos (2003): Fumigant toxicity of three essential oils on the eggs of *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae). Available online at sciencedirect.com

21.Paulraj, Koundal,(2002) Plant protease inhibitors in control of pytophagous insects, April 15.

22. P.G.Fenemore, Alka Prakash,(1961), Applied Entomology, Wiley Eastern LTD.

23.Raksha Malakar, (2000), Glandular trichomes of Solanum berthaultii and its hybrids with potato deter oviposotion and impair growthof poatato tuber moth, Entomologia Experimentalis et Applocata, 94(3):249-257.

24.Regnault-Roger C, (1997). The potential of botanical essential oils for insect pest control. Integr. Pest Manage. Rev. 2,25–34.

25. Renolds, J.F.,(1989). Martindale, The Extra Phamacopoeia. The Pharceutical Press. London. pp: 132.

26.Schmutterer, H., (1995). The Tree and its Characteristics. *In* Schmutterer, H. (Eds.). The Neem Tree, VCH Weinheim, Germany, pp: 1-34.

27.Schmutterer, H., (1988). Potential of azadirachtincontaining pesticides for integrated pest control in developing and industrialized countries. J Insect Physiol 34: 713-719.

28.Sharma R.N (1995): Modern approaches to the development and utilization of plant products for pest management: A holistic review. In 'Cultivation and Utilization of Medicinal Plants'. Eds.K.K.Atal and B.M.Kapur. Jammu Tawi.

29.Sharma R.N., Deshpande S.G., Joseph M. (1994): Toxicity of natural essential oils against Aedes aegypti and Culex fatigans., *Geobios*, 21(3), 162-165.

30.Sharma, S.K., V.K. Dua and V.P. Sharma, (1995). Field studies on the mosquito repellent action of neem oil. Southeast Asian J. Trop. Med. Public Health., 26: 180-2.

31.Sorensen, M.,(1996). Yam bean. Promoting the conservation and use of underutilized and neglected crops. 2.International Plant Genetic Resource Institute: Rome.

32. Stamford, N.P., D.S. Ferraz, A.D.S. Freitas and C.E.R.S.Santos, (2002). Effect of sulphur inoculated with *Thiobacillus* on saline soils amendment and growth of cowpea and yam bean legumes. J. Agric. Sci., Cambridge, 139: 1-7.

33.Subra, R.,(1980). Biology and control of *Culex quinquefasciatus* Say (Diptera: Culicidae). Eight special reference to Africa. WHO Mimeographed Document Series, No. WHO/VBC/80.781. pp: 44.

34.Tsao R and Coats J R (1995): Starting from nature to make better insecticides, *Chemtech.*25, 23-38.

35.Whittakar R.H. and Fenny R.P.(1971): Allelochemicals: Chemical Interaction between species, *Science* 171, 757-770.

Name of the Student: Mr. Rahul Krishna Pawar

Name and Signature of Co-guide: Dr.(Mrs). Rashmi Morey

Name and Signature of Guide: Dr. Abhay J. Khandagle